arXiv:astro-ph/0609446v1 15 Sep 2006

FULL TITLE
*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***
NAMES OF EDITORS

Optical & Infrared Observations of Stellar Mass Loss in Globular Cluster Red Giants

Iain McDonald[1], Jacco Th. van Loon[1]

[1]*Astrophysics Group, Lennard Jones Laboratories, EPSAM, Keele University, Staffordshire, ST5 5BG, UK*

Abstract. We are examining mass loss from globular clusters giant stars, focussing on metallicity dependance. We present three sets of observations: TIMMI-2 mid-IR spectra of 47 Tuc, UVES high-resolution optical spectra of several clusters, and an infrared atlas of ω Cen using the Spitzer Space Telescope.

1. Introduction

Mass loss from giants is of great importance to stellar evolution — low-mass stars lose up to 0.2 $M_\odot$ on the giant branch. Metal-poor globulars giants can probe metallicity dependances of this mass loss, exposing the Universe's chemical enrichment history, and revealing how low-metallicity stars can produce large quantities of dust (*e.g.* Boyer et al. 2006).

2. Mid-Infrared Spectroscopy in 47 Tucanae

Eight infrared-excessive giant branch stars in 47 Tuc were observed in the mid-infrared with the ESO La Silla 3.6m and TIMMI-2 spectrograph. Clear silicate emission is present in V1, corresponding to a mass loss rate of 10^{-6} $M_\odot$ yr^{-1} (estimated using the DUSTY code — Ivezić, Nenkova & Elizur (1999)). This work was published as van Loon et al. (2006).

3. Optical Spectroscopy of Cluster Giants

Optical VLT/UVES spectra were taken of 47 giant stars in six clusters, shown in Fig. 1, in order to analyse differences between IR-excessive and normal stars at a range of metallicities. At $R > 100{,}000$, these represent some of the highest resolution data ever taken of globular cluster giants.

The Hα and mid-IR Ca II triplet lines exhibit evidence for mass loss: core shifts of several km s^{-1} exist in many stars, as does substantial emission. We are in the process of qualitatively analysing the Hα lines using the SEI code (Lamers, Cerruti-Sola & Perinotto 1987) and various other computational methods. We will then perform an abundance analysis and attempt to model the chromospheric emission.



Figure 1. [Left] UVES spectra, sorted by spectral type and normalised to their respective 7000 Å fluxes. The spectra have been smoothed for clarity. The central inset details the Hα line. [Right] Example profiles from the inset, showing examples of core asymmetries, emission and emission asymmetries and pulsation-shocked emission.

4. Spitzer Space Telescope Infrared Atlas of ω Centauri

Spitzer imaging of ω Cen totalling 18.4 hours was taken in six bands from 3.6–70 μm. We will produce the first multi-wavelength atlas of ω Cen in the infrared. We can then use this to find dust-enshrouded objects in the cluster, and measure their mass loss rates and other characteristics.

Acknowledgments. We are thankful to JPL/NASA and ESO for the use of their facilities. Iain McDonald is supported by a PPARC studentship.

References

Boyer, M. L., Woodward, C. E., van Loon, J. Th., Gordon, K., Evans, A., Gehrz, R. D., Helton, L. A., Polomski, E. F. 2006, AJ, 132, 1415

Ivezić, Z., Nenkova, M., & Elizur, M. 1999, User manual for DUSTY, University of Kentucky Internal Report

Lamers, H. J. G. L. M., Cerruti-Sola, M., & Perinotto, M. 1987, Ap. J., 314, 726

van Loon, J. Th., McDonald, I., Oliveira, J. M., Evans, A., Boyer, M .L., Gehrz, R. D., Polomski, E., Woodward, C. E. 2006, A&A, 450, 339